Volume 84 No. 1

Ecological MONOGRAPHS

A PUBLICATION OF THE ECOLOGICAL SOCIETY OF AMERICA



Articles

A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche partitioning

Phylogenetic beta diversity, similarity, and differentiation measures based on Hill numbers

Stochastic models reveal conditions for cyclic dominance in sockeye salmon populations

Ecological Monographs

VOL. 84 • NO. 1 • FEBRUARY 2014 ISSN 0012-9615

CONTENTS

Editorial

A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche partitioning • D. LEE TAYLOR, TERESA N. HOLLINGSWORTH, JACK W. MCFARLAND, NIALL J. LENNON, CHAD NUSBAUM, AND ROGER W. RUESS

Phylogenetic beta diversity, similarity, and differentiation measures based on Hill numbers

CHUN-HUO CHIU, LOU JOST, AND ANNE CHAO

Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species

• ANNE CHAO, NICHOLAS J. GOTELLI, T. C. HSIEH, ELIZABETH L. SANDER, K. H. MA, ROBERT K. COLWELL, AND AARON M. ELLISON

Stochastic models reveal conditions for cyclic dominance in sockeye salmon populations • J. WILSON WHITE, LOUIS W. BOTSFORD, ALAN HASTINGS, AND MATTHEW D. HOLLAND

Intercontinental comparison of fish ecomorphology: null model tests of community assembly at the patch scale in rivers

CARMEN G. MONTAÑA, KIRK O. WINEMILLER, AND ANDREW SUTTON

Carbon transit through degradation networks

DAVID C. FORNEY AND DANIEL H. ROTHMAN

How climate extremes—not means—define a species' geographic range boundary via a demographic

• HEATHER J. LYNCH, MARC RHAINDS, JUSTIN M. CALABRESE, STEPHEN CANTRELL, CHRIS COSNER, AND WILLIAM F. FAGAN

Responses of a tundra system to warming using SCAMPS: a stoichiometrically coupled, acclimating microbe-plant-soil model

• SEETA A. SISTLA, EDWARD B. RASTETTER, AND JOSHUA P. SCHIMEL

Appendices and Supplements are available online: www.esapubs.org/archive

Instructions to Authors Available online: www.esapubs.org/esapubs/AuthorInstructions

COVER PHOTO: Mushrooms of a Cortinarius species (possibly C. favrei) from Alaska, USA. Cortinarius was the most abundant genus found in boreal forest soils by Taylor et al., with strong niche-partitioning among species (see pp. 3-20). Photo credit: Ima Timling.